

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 September 2001 (27.09.2001)

PCT

(10) International Publication Number
WO 01/70995 A1

(51) International Patent Classification⁷: C12N 15/53,
15/77, C12P 13/08

(21) International Application Number: PCT/EP00/06303

(22) International Filing Date: 5 July 2000 (05.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/531,269 20 March 2000 (20.03.2000) US

(71) Applicants: DEGUSSA AG [DE/DE]; Bennigsenplatz
1, 40474 Düsseldorf (DE). FORSCHUNGSZENTRUM
JÜLICH GMBH [DE/DE]; 52425 Jülich (DE). NA-
TIONAL UNIVERSITY OF IRELAND [IE/IE];
Galway (IE).

(72) Inventors: BURKE, Kevin; 5, Greenfield Road, New-
castle, Galway, County Galway (IE). SAHM, Hermann;

Wendelinusstrasse 71, 52428 Jülich (DE). EGGELING,
Lothar; Elsenkamp 6, 52428 Jülich (DE). MORITZ,
Bernd; Schattbachstrasse 1, 44801 Bochum (DE). DUNI-
CAN, L., K. (deceased). MCCORMACK, Ashling;
Moate Road, Athlone, County Westmeath (IE). STAPEL-
TON, Cliona; 27, Railway View, Roscrea, County
Tipperary (IE). MÖCKEL, Bettina; Benrodestrasse 35,
40597 Düsseldorf (DE). THIERBACH, Georg; Gunst-
strasse 21, 33613 Bielefeld (DE).

(81) Designated States (*national*): AU, BR, CA, CN, HU, ID,
JP, KR, MX, PL, RU, SK, UA, ZA.

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

Published:
— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: PROCESS FOR THE FERMENTATIVE PREPARATION OF L-AMINO ACIDS WITH AMPLIFICATION OF THE
ZWF GENE

(57) Abstract: The invention relates to a process for the preparation of L-amino acids by fermentation of coryneform bacteria, which
comprises carrying out the following steps: a) fermentation of the desired L-amino acid-producing bacteria in which at least the zwf
gene is amplified, b) concentration of the L-amino acid in the medium or in the cells of the bacteria and c) isolation of the L-amino
acid produced.



WO 01/70995 A1

**Process for the fermentative preparation of L-amino acids
with amplification of the zwf gene**

The invention relates to a process for the fermentative preparation of L-amino acids, in particular L-lysine, L-
5 threonine and L-tryptophan, using coryneform bacteria in which at least the zwf gene is amplified.

Prior art

L-Amino acids are used in animal nutrition, in human medicine and in the pharmaceuticals industry.

10 It is known that amino acids are prepared by fermentation of strains of coryneform bacteria, in particular Corynebacterium glutamicum. Because of its great importance, work is constantly being undertaken to improve the preparation process. Improvements to the process can
15 relate to fermentation measures, such as e. g. stirring and supply of oxygen, or the composition of the nutrient media, such as e. g. the sugar concentration during the fermentation, or the working up to the product form by
20 e. g. ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to
25 antimetabolites, such as e. g. the threonine analogue α -amino- β -hydroxyvaleric acid (AHV), or are auxotrophic for metabolites of regulatory importance and produce L-amino acids such as e. g. threonine are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of
30 Corynebacterium glutamicum strains which produce L-amino acids.

Object of the invention

The inventors had the object of providing new improved processes for the fermentative preparation of L-amino acids with coryneform bacteria.

5 Description of the invention

L-Amino acids are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and especially in animal nutrition. There is therefore a general interest in providing new improved processes for
10 the preparation of amino acids.

The invention provides a process for the fermentative preparation of L-amino acids, in particular L-lysine, L-threonine and L-tryptophan, using coryneform bacteria in which the nucleotide sequence which codes for the Zwf
15 protein (zwf gene) is amplified, in particular over-expressed.

The strains employed preferably already produce L-amino acids before amplification of the zwf gene.

Preferred embodiments are to be found in the claims.

20 The term "amplification" in this connection describes the increase in the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or
25 using a gene which codes for a corresponding enzyme (protein) having a high activity, and optionally combining these measures.

The microorganisms which the present invention provides can prepare L-amino acids from glucose, sucrose, lactose,
30 fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They are representatives of

coryneform bacteria, in particular of the genus
Corynebacterium. Of the genus Corynebacterium, there may be
mentioned in particular the species Corynebacterium
glutamicum, which is known among specialists for its
5 ability to produce L-amino acids.

Suitable strains of the genus Corynebacterium, in
particular of the species Corynebacterium glutamicum, are,
for example, the known wild-type strains

10 Corynebacterium glutamicum ATCC13032
Corynebacterium acetoglutamicum ATCC15806
Corynebacterium acetoacidophilum ATCC13870
Corynebacterium thermoaminogenes FERM BP-1539
Brevibacterium flavum ATCC14067
Brevibacterium lactofermentum ATCC13869
15 Brevibacterium divaricatum ATCC14020

and L-amino acid-producing mutants prepared therefrom,

such as, for example, the L-threonine-producing strains

Corynebacterium glutamicum ATCC21649
Brevibacterium flavum BB69
20 Brevibacterium flavum DSM5399
Brevibacterium lactofermentum FERM-BP 269
Brevibacterium lactofermentum TBB-10

and such as, for example, the L-isoleucine-producing
strains

25 Corynebacterium glutamicum ATCC 14309
Corynebacterium glutamicum ATCC 14310
Corynebacterium glutamicum ATCC 14311
Corynebacterium glutamicum ATCC 15168
Corynebacterium ammoniagenes ATCC 6871

30 and such as, for example, the L-tryptophan-producing
strains

Corynebacterium glutamicum ATCC21850 and
Corynebacterium glutamicum KY9218 (pKW9901)

and such as, for example, the L-lysine-producing strains

Corynebacterium glutamicum FERM-P 1709

Brevibacterium flavum FERM-P 1708

Brevibacterium lactofermentum FERM-P 1712

5 *Corynebacterium glutamicum* FERM-P 6463

Corynebacterium glutamicum FERM-P 6464

Corynebacterium glutamicum ATCC13032

Corynebacterium glutamicum DM58-1

Corynebacterium glutamicum DSM12866.

10 It has been found that coryneform bacteria produce L-amino acids, in particular L-lysine, L-threonine and L-tryptophan, in an improved manner after over-expression of the *zwf* gene which codes for the Zwf protein.

Alleles of the *zwf* gene which result from the degeneracy of
15 the genetic code or due to sense mutations of neutral function can furthermore be used.

JP-A-09224661 discloses the nucleotide sequence of the glucose 6-phosphate dehydrogenase gene, called *zwf*, of *Brevibacterium flavum* MJ-223 (FERM BP-1497). JP-A-09224661
20 describes the N-terminal amino acid sequence of the Zwf polypeptide as Met Val Ile Phe Gly Val Thr Gly Asp Leu Ala Arg Lys Lys Leu.

However, it has not been possible to confirm this. Instead, the following N-terminal amino acid sequence has been
25 found: Val Ser Thr Asn Thr Thr Pro Ser Ser Trp Thr Asn Pro Leu Arg Asp. The valyl radical in the N-position can be split off in the context of post-translational modification, and Ser Thr Asn Thr Thr Pro Ser Ser Trp Thr Asn Pro Leu Arg Asp is then obtained as the N-terminal
30 amino acid sequence.

To achieve an amplification (e. g. over-expression), the number of copies of the corresponding genes is increased, or the promoter and regulation region or the ribosome

binding site upstream of the structural gene is mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally possible to increase the expression in the course of fermentative L-amino acid formation. The expression is likewise improved by measures to prolong the life of the m-RNA. Furthermore, the enzyme activity is also increased by preventing the degradation of the enzyme protein. The genes or gene constructs are either present here in plasmids with a varying number of copies, or are integrated and amplified in the chromosome. Alternatively, an over-expression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

Instructions in this context can be found by the expert, inter alia, in Martin et al. (Bio/Technology 5, 137-146 (1987)), in Guerrero et al. (Gene 138, 35-41 (1994)), Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)), in Eikmanns et al. (Gene 102, 93-98 (1991)), in European Patent Specification EPS 0 472 869, in US Patent 4,601,893, in Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)), in Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)), in LaBarre et al. (Journal of Bacteriology 175, 1001-1007 (1993)), in Patent Application WO 96/15246, in Malumbres et al. (Gene 134, 15-24 (1993)), in Japanese Laid-Open Specification JP-A-10-229891, in Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998)) and in known textbooks of genetics and molecular biology.

By way of example, the Zwf protein was over-expressed with the aid of a plasmid. The E. coli - C. glutamicum shuttle vector pEC-T18mob2 shown in Figure 1 was used for this. After incorporation of the zwf gene into the KpnI/SalI cleavage site of pEC-T18mob2, the plasmid pEC-T18mob2zwf shown in Figure 2 was formed.

Other plasmid vectors which are capable of replication in *C. glutamicum*, such as e.g. pEKEx1 (Eikmanns et al., Gene 102:93-98 (1991)) or pZ8-1 (EP-B- 0 375 889), can be used in the same way.

- 5 In addition, it may be advantageous for the production of L-amino acids to amplify one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the pentose phosphate pathway or of amino acid export, in addition to amplification of the *zwf* gene.
- 10 Thus, for example, in particular for the preparation of L-threonine, one or more genes chosen from the group consisting of:
- the *hom* gene which codes for homoserine dehydrogenase (Peoples et al., Molecular Microbiology 2, 63-72 (1988))
15 or the *hom^{dr}* allele which codes for a "feed back resistant" homoserine dehydrogenase (Archer et al., Gene 107, 53-59 (1991)),
 - the *gap* gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns et al., Journal of Bacteriology
20 174: 6076-6086 (1992)),
 - the *pyc* gene which codes for pyruvate carboxylase (Peters-Wendisch et al., Microbiology 144: 915-927 (1998)),
 - the *mgo* gene which codes for malate:quinone
25 oxidoreductase (Molenaar et al., European Journal of Biochemistry 254, 395-403 (1998)),
 - the *tkt* gene which codes for transketolase (accession number AB023377 of the European Molecular Biologies Laboratories databank (EMBL, Heidelberg, Germany)),

- the gnd gene which codes for 6-phosphogluconate dehydrogenase (JP-A-9-224662),
- the thrE gene which codes for threonine export (DE 199 41 478.5; DSM 12840),

5 • the zwf gene (DE 199 59 328.0; DSM 13115),

- the eno gene which codes for enolase (DE: 199 41 478.5)

can be amplified, in particular over-expressed, at the same time.

Thus, for example, in particular for the preparation of L-lysine, one or more genes chosen from the group consisting of

- the dapA gene which codes for dihydrodipicolinate synthase (EP-B 0 197 335),
- the lysC gene which codes for a feed back resistant aspartate kinase (Kalinowski et al. (1990), Molecular and General Genetics 224: 317-324),
- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- 20 • the pyc gene which codes for pyruvate carboxylase (DE-A-198 31 609),
- the tkt gene which codes for transketolase (accession number AB023377 of the European Molecular Biologies Laboratories databank (EMBL, Heidelberg, Germany)), «
- 25 • the gnd gene which codes for 6-phosphogluconate dehydrogenase (JP-A-9-224662),

- the lysE gene which codes for lysine export (DE-A-195 48 222),
- the zwf gene (DE 199 59 328.0; DSM 13115),
- the eno gene which codes for enolase (DE 199 47 791.4)

5 can be amplified, in particular over-expressed, at the same time.

It may furthermore be advantageous for the production of L-amino acids at the same time to attenuate one of the genes chosen from the group consisting of

- 10 • the pck gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1; DSM 13047),
- the pgf gene which codes for glucose 6-phosphate isomerase (US 09/396,478, DSM 12969),
- the poxB gene which codes for pyruvate oxidase
- 15 (DE 199 51 975.7; DSM 13114),
- the zwf2 gene (DE: 199 59 327.2; DSM 13113)

in addition to the amplification of the zwf gene.

In addition to over-expression of the Zwf protein, it may furthermore be advantageous for the production of L-amino
20 acids to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The microorganisms prepared according to the invention can
25 be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of L-amino acid production. A summary of known culture methods is described in the textbook by Chmiel

(Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen [Bioreactors and Peripheral Equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular microorganisms in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981). Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as e. g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e. g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e. g. glycerol and ethanol, and organic acids, such as e. g. acetic acid, can be used as the source of carbon. These substance can be used individually or as a mixture. Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulphate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture. Potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as e. g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the above-mentioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances

mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium
5 hydroxide, ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective
10 action, e.g. antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and
15 preferably 25°C to 40°C. Culturing is continued until a maximum of L-amino acid has formed. This target is usually reached within 10 hours to 160 hours.

The analysis of L-amino acids can be carried out by anion exchange chromatography with subsequent ninhydrin
20 derivatization, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190), or it can take place by reversed phase HPLC as described by Lindroth et al. (Analytical Chemistry (1979) 51: 1167-1174).

The following microorganism has been deposited at the
25 Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty:

Escherichia coli K-12 DH5 α /pEC-T18mob2 as DSM 13244

The following figures are attached:

- Figure 1: Map of the plasmid pEC-T18mob2
- Figure 2: Map of the plasmid pEC-T18mob2zwf
- Figure 3: Map of the plasmid pAMC1
- 5 • Figure 4: Map of the plasmid pMC1
- Figure 5: Map of the plasmid pCR2.1poxBint

The base pair numbers stated are approx. values obtained in the context of reproducibility.

Re Figure 1 and 2:

- 10 The abbreviations used have the following meaning:

	Tet:	Resistance gene for tetracycline
	oriV:	Plasmid-coded replication origin of <i>E. coli</i>
	RP4mob:	mob region for mobilizing the plasmid
	rep:	Plasmid-coded replication origin from
15		<i>C. glutamicum</i> plasmid pGA1
	per:	Gene for controlling the number of copies
		from pGA1
	lacZ-alpha:	lacZ α gene fragment (N-terminus) of the
		β -galactosidase gene
20	lacZalpha':	5'-Terminus of the lacZ α gene fragment
	'lacZalpha:	3'-Terminus of the lacZ α gene fragment

Re Figure 3 and 4:

The abbreviations used have the following meaning:

	Neo r :	Neomycin/kanamycin resistance
25	ColE1 ori:	Replication origin of the plasmid ColE1
	CMV:	Cytomegalovirus promoter
	lacP:	Lactose promoter
	pgi:	Phosphoglucose isomerase gene
	lacZ:	Part of the β -galactosidase gene
30	SV40 3' splice	3' splice site of Simian virus 40

SV40 polyA: Polyadenylation site of Simian virus
40
fl(-)ori: Replication origin of the filamentous
phage fl
5 SV40 ori: Replication origin of Simian virus
40
kan r: Kanamycin resistance
pgi insert: Internal fragment of the pgi gene
ori: Replication origin of the plasmid pBGS8

10 Re Figure 5:

The abbreviations used have the following meaning:

ColE1 ori: Replication origin of the plasmid ColE1
lacZ: Cloning relict of the lacZ α gene fragment
fl ori: Replication origin of phage fl
15 KmR: Kanamycin resistance
ApR: Ampicillin resistance
poxBint: Internal fragment of the poxB gene

The meaning of the abbreviations for the various
restriction enzymes (e. g. BamHI, EcoRI etc.) are known from
20 the prior art and are summarized, for example, by Kessler
and Höltnke (Gene 47, 1-153 (1986)) or Roberts et al.
(Nucleic Acids Research 27, 312-313 (1999)).

Examples

The following examples will further illustrate this invention. The molecular biology techniques, e.g. plasmid DNA isolation, restriction enzyme treatment, ligations, standard transformations of *Escherichia coli* etc. used are, (unless stated otherwise), described by Sambrook et al., (Molecular Cloning. A Laboratory Manual (1989) Cold Spring Harbour Laboratories, USA).

Example 1

10 Expression of the Zwf protein

1.1 Preparation of the plasmid pEC-T18mob2

The *E. coli* - *C. glutamicum* shuttle vector pEC-T18mob2 was constructed according to the prior art. The vector contains the replication region rep of the plasmid pGA1 including the replication effector per (US-A- 5,175,108; Nesvera et al., Journal of Bacteriology 179, 1525-1532 (1997)), the tetracycline resistance-imparting tetA(2) gene of the plasmid pAG1 (US-A- 5,158,891; gene library entry at the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA) with accession number AF121000), the replication region oriV of the plasmid pMB1 (Sutcliffe, Cold Spring Harbor Symposium on Quantitative Biology 43, 77-90 (1979)), the lacZ α gene fragment including the lac promoter and a multiple cloning site (mcs) (Norrander et al. Gene 26, 101-106 (1983)) and the mob region of the plasmid RP4 (Simon et al., (1983) Bio/Technology 1:784-791). The vector constructed was transformed in the *E. coli* strain DH5 α (Brown (ed.) Molecular Biology Labfax, BIOS Scientific Publishers, Oxford, UK, 1991). Selection for plasmid-carrying cells was made by plating out the transformation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which had been supplemented with 5 mg/l tetracycline. Plasmid DNA was

isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme EcoRI and HindIII and subsequent agarose gel electrophoresis (0.8%).

- 5 The plasmid was called pEC-Tl8mob2 and is shown in Figure 1. It is deposited in the form of the strain Escherichia coli K-12 strain DH5 α pEC-Tl8mob2 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, 10 Braunschweig, Germany) as DSM 13244.

1.2 Preparation of the plasmid pEC-Tl8mob2zwf

The gene from Corynebacterium glutamicum ATCC13032 was first amplified by a polymerase chain reaction (PCR) by means of the following oligonucleotide primer:

- 15 zwf-forward:

5'-TCG ACG CGG TTC TGG AGC AG-3'

zwf-reverse:

5'-CTA AAT TAT GGC CTG CGC CAG-3'

- The PCR reaction was carried out in 30 cycles in the 20 presence of 200 μ M deoxynucleotide triphosphates (dATP, dCTP, dGTP, dTTP), in each case 1 μ M of the corresponding oligonucleotide, 100 ng chromosomal DNA from Corynebacterium glutamicum ATCC13032, 1/10 volume 10-fold reaction buffer and 2.6 units of a heat-stable Taq-/Pwo-DNA 25 polymerase mixture (Expand High Fidelity PCR System from Roche Diagnostics, Mannheim, Germany) in a Thermocycler (PTC-100, MJ Research, Inc., Watertown, USA) under the following conditions: 94°C for 30 seconds, 64°C for 1 minute and 68°C for 3 minutes.

- 30 The amplified fragment about 1.8 kb in size was subsequently ligated with the aid of the SureClone Ligation Kit (Amersham Pharmacia Biotech, Uppsala, Sweden) into the SmaI cleavage site of the vector pUC18 in accordance with

the manufacturer's instructions. The E. coli strain DH5 α mcr (Grant et al., Proceedings of the National Academy of Sciences of the United States of America USA (1990) 87: 4645-4649) was transformed with the entire ligation batch.

5 Transformants were identified with the aid of their carbenicillin resistance on LB-agar plates containing 50 μ g/mL carbenicillin. The plasmids were prepared from 7 of the transformants and checked for the presence of the 1.8 kb PCR fragment as an insert by restriction analysis.

10 The recombinant plasmid formed in this way is called pUC18zwf in the following.

For construction of pEC-T18mob2zwf, pUC18zwf was digested with KpnI and SalI, and the product was isolated with the aid of the NucleoSpin Extraction Kit from Macherey-Nagel

15 (Düren, Germany) in accordance with the manufacturer's instructions and then ligated with the vector pEC-T18mob2, which had also been cleaved with KpnI and SalI and dephosphorylated. The E. coli strain DH5 α mcr (Grant et al., Proceedings of the National Academy of Sciences of the

20 United States of America USA (1990) 87: 4645-4649) was transformed with the entire ligation batch. Transformants were identified with the aid of their tetracycline resistance on LB-agar plates containing 5 μ g/mL tetracycline. The plasmids were prepared from 12 of the

25 transformants and checked for the presence of the 1.8 kb PCR fragment as an insert by restriction analysis. One of the recombinant plasmids isolated in this manner was called pEC-T18mob2zwf (Figure 2).

Example 2

30 Preparation of amino acid producers with an amplified zwf gene

The L-lysine-producing strain *Corynebacterium glutamicum* DSM5715 is described in EP-B-0435132 and the L-threonine-producing strain *Brevibacterium flavum* DSM5399 is described

in EP-B-0385940. Both strains are deposited at the Deutsche Sammlung für Mikroorganismen und Zellkulturen [German Collection of Microorganisms and Cell Cultures] in Braunschweig (Germany) in accordance with the Budapest Treaty.

2.1 Preparation of the strains DSM5715/pEC-T18mob2zwf and DSM5399/pEC-T18mob2zwf

The strains DSM5715 and DSM5399 were transformed with the plasmid pEC-T18mob2zwf using the electroporation method described by Liebl et al., (FEMS Microbiology Letters, 53:299-303 (1989)) Selection of the transformants took place on LBHIS agar comprising 18.5 g/l brain-heart infusion broth, 0.5 M sorbitol, 5 g/l Bacto-tryptone, 2.5 g/l Bacto-yeast extract, 5 g/l NaCl and 18 g/l Bacto-agar, which had been supplemented with 5 mg/l tetracycline. Incubation was carried out for 2 days at 33°C.

Plasmid DNA was isolated in each case from a transformant by conventional methods (Peters-Wendisch et al., 1998, Microbiology 144, 915 -927), cleaved with the restriction endonucleases XbaI and KpnI, and the plasmid was checked by subsequent agarose gel electrophoresis. The strains obtained in this way were called DSM5715/pEC-T18mob2zwf and DSM5399/pEC-T18mob2zwf.

2.2 Preparation of L-threonine

The *C. glutamicum* strain DSM5399/pEC-T18mob2zwf obtained in Example 2.1 was cultured in a nutrient medium suitable for the production of threonine and the threonine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with tetracycline (5 mg/l)) for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml

medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture.

Medium Cg III

NaCl	2.5 g/l
------	---------

Bacto-Peptide	10 g/l
---------------	--------

Bacto-Yeast extract	10 g/l
---------------------	--------

Glucose (autoclaved separately) 2% (w/v)

The pH was brought to pH 7.4

Tetracycline (5 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking
5 machine. A main culture was seeded from this preculture such that the initial OD (660nm) of the main culture was 0.1. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50 g/l
 (NH ₄) ₂ SO ₄	 25 g/l
KH ₂ PO ₄	0.1 g/l
MgSO ₄ * 7 H ₂ O	1.0 g/l
CaCl ₂ * 2 H ₂ O	10 mg/l
FeSO ₄ * 7 H ₂ O	10 mg/l
MnSO ₄ * H ₂ O	5.0 mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
L-Leucine (sterile-filtered)	0.1 g/l
CaCO ₃	25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO₃,
5 autoclaved in the dry state.

Culturing is carried out in a 10 ml volume in a 100 ml conical flask with baffles. Tetracycline (5 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of threonine formed was determined with an amino acid analyzer from Eppendorf-
5 BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in Table 1.

Table 1

Strain	OD	L-Threonin g/l
DSM5399	12.3	0.74
DSM5399/pEC-T18mob2zwf	10.2	1.0

10 2.3 Preparation of L-lysine

The *C. glutamicum* strain DSM5715/pEC-T18mob2zwf obtained in Example 2.1 was cultured in a nutrient medium suitable for the production of lysine and the lysine content in the culture supernatant was determined.

- 15 For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with tetracycline (5 mg/l)) for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium
20 CgIII was used as the medium for the preculture.

Medium Cg III

NaCl	2.5 g/l
Bacto-Peptide	10 g/l

Bacto-Yeast extract 10 g/l

Glucose (autoclaved separately) 2% (w/v)

The pH was brought to pH 7.4

Tetracycline (5 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660nm) of the main culture was

5 0.1. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor) 5 g/l

MOPS (morpholinopropanesulfonic acid) 20 g/l

Glucose (autoclaved separately) 58 g/l

(NH₄)₂SO₄ 25 g/l

KH₂PO₄ 0.1 g/l

MgSO₄ * 7 H₂O 1.0 g/l

CaCl₂ * 2 H₂O 10 mg/l

FeSO₄ * 7 H₂O 10 mg/l

MnSO₄ * H₂O 5.0mg/l

Biotin (sterile-filtered) 0.3 mg/l

Thiamine * HCl (sterile-filtered) 0.2 mg/l

L-Leucine (sterile-filtered) 0.1 g/l

CaCO₃ 25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO₃ autoclaved in the dry state.

- 5 Culturing is carried out in a 10 ml volume in a 100 ml conical flask with baffles. Tetracycline (5 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

- 10 After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

- 15 The result of the experiment is shown in Table 2.

Table 2

Strain	OD	L-Lysine HCl g/l
DSM5715	10.8	16.0
DSM5715/pEC-T18mob2zwf	7.2	17.1

Example 3

- Construction of a gene library of *Corynebacterium*
20 glutamicum strain AS019

- A DNA library of *Corynebacterium glutamicum* strain AS019 (Yoshihama et al., Journal of Bacteriology 162, 591-597 (1985)) was constructed using λ Zap Express™ system, (Short et al., (1988) Nucleic Acids Research, 16: 7583-
25 7600), as described by O'Donohue (O'Donohue, M. (1997). The

Cloning and Molecular Analysis of Four Common Aromatic Amino Acid Biosynthetic Genes from *Corynebacterium glutamicum*. Ph.D. Thesis, National University of Ireland, Galway). λ Zap Express™ kit was purchased from Stratagene (Stratagene, 11011 North Torrey Pines Rd., La Jolla, California 92037) and used according to the manufacturers instructions. AS019-DNA was digested with restriction enzyme *Sau3A* and ligated to *Bam*HI treated and dephosphorylated λ Zap Express™ arms.

10 Example 4

Cloning and sequencing of the *pgi* gene

1. Cloning

Escherichia coli strain DF1311, carrying mutations in the *pgi* and *pgl* genes as described by Kupor and Fraenkel, (Journal of Bacteriology 100: 1296-1301 (1969)), was transformed with approx. 500 ng of the AS019 λ Zap Express™ plasmid library described in Example 3. Selection for transformants was made on M9 minimal media, (Sambrook et al., (1989). Molecular Cloning. A Laboratory Manual Cold Spring Harbour Laboratories, USA), containing kanamycin at a concentration of 50 mg/l and incubation at 37°C for 48 hours. Plasmid DNA was isolated from one transformant according to Birnboim and Doly (Nucleic Acids Research 7: 1513-1523 (1979)) and designated pAMC1 (Figure 3).

25 2. Sequencing

For sequence analysis of the cloned insert of pAMC1 the method of Sanger et al. (Proceedings of the National Academy of Sciences USA 74,5463-5467 (1977)) was applied using primers differentially labelled with a coloured fluorescent tag. It was carried out using the ABI prism 310 genetic analyzer from Perkin Elmer Applied Biosystems, (Perkin Elmer Corporation, Norwalk, Connecticut, U.S.A), and the ABI prism Big Dye™ Terminator Cycle Sequencing Ready Reaction kit also from Perkin Elmer.

Initial sequence analysis was carried out using the universal forward and M13 reverse primers obtained from Pharmacia Biotech (St. Albans, Herts, AL1 3AW, UK):

Universal forward primer: GTA ATA CGA CTC ACT ATA GGG C

5 M13 reverse primer: GGA AAC AGC TAT GAC CAT G

Internal primers were subsequently designed from the sequence obtained which allowed the entire *pgi* gene to be deduced. The sequence of the internal primers is as follows:

10 Internal primer 1: GGA AAC AGG GGA GCC GTC

Internal primer 2: TGC TGA GAT ACC AGC GGT

Sequence obtained was then analyzed using the DNA Strider programme, (Marck, (1988). Nucleic Acids Research 16: 1829-1836), version 1.0 on an Apple Macintosh computer. This
15 program allowed for analyses such as restriction site usage, open reading frame analysis and codon usage determination. Searches between DNA sequence obtained and those in EMBL and Genbank databases were achieved using the BLAST programme, (Altschul et al., (1997). Nucleic Acids
20 Research, 25: 3389-3402). DNA and protein sequences were aligned using the Clustal V and Clustal W programs (Higgins and Sharp, 1988 Gene 73: 237-244).

The sequence thus obtained is shown in SEQ ID NO 1. The analysis of the nucleotide sequence obtained revealed an
25 open reading frame of 1650 base pairs which was designated as *pgi* gene. It codes for a protein of 550 amino acids shown in SEQ ID NO 2.

Example 5

Preparation of an integration vector for integration
30 mutagenesis of the *pgi* gene

An internal segment of the *pgi* gene was amplified by polymerase chain reaction (PCR) using genomic DNA isolated from *Corynebacterium glutamicum* AS019, (Heery and Dunican,

(1993) Applied and Environmental Microbiology 59: 791-799), as template. The pgi primers used were:

fwd. Primer: ATG GAR WCC AAY GGH AA

rev. Primer: YTC CAC GCC CCA YTG RTC

5 with R=A+G; Y=C+T; W=A+T; H=A+T+C.

PCR Parameters were as follows: 35 cycles

94°C for 1 min.

47°C for 1 min.

72°C for 30 sec.

10 1.5 mM MgCl₂

approx. 150-200 ng DNA template.

The PCR product obtained was cloned into the commercially available pGEM-T vector received from Promega Corp., (Promega UK, Southampton.) using strain E. coli JM109, (Yanisch-Perron et al., 1985. Gene, 33: 103-119), as a
15 host. The sequence of the PCR product is shown as SEQ ID NO 3. The cloned insert was then excised as an EcoRI fragment and ligated to plasmid pBGS8 (Spratt et al., Gene 41: 337-342 (1986)) pretreated with EcoRI. The restriction enzymes
20 used were obtained from Boehringer Mannheim UK Ltd., (Bell Lane, Lewes East Sussex BN7 1LG, UK.) and used according to manufacturers instructions. E. coli JM109 was then transformed with this ligation mixture and electrotransformants were selected on Luria agar
25 supplemented with IPTG (isopropyl- β -D-thiogalactopyranoside), XGAL (5-bromo-4-chloro-3-indolyl-D-galactopyranoside) and kanamycin at a concentration of 1 mM, 0.02% and 50 mg/l respectively. Agar plates were incubated for twelve hours at 37°C. Plasmid DNA was
30 isolated from one transformant, characterized by restriction enzyme analysis using EcoRI, BamHI and SalI designated pMC1 (Figure 4).

Plasmid pMC1 was deposited in the form of Escherichia coli strain DH5a/pMC1 at the Deutsche Sammlung für

Mikroorganismen und Zellkulturen (DSMZ, Braunschweig, Germany) as DSM 12969 according to the Budapest treaty.

Example 6

Integration mutagenesis of the *pgi* gene in the lysine
5 producer DSM 5715

The vector pMC1 mentioned in Example 5 was electroporated by the electroporation method of Tauch et al. (FEMS Microbiological Letters, 123:343-347 (1994)) in *Corynebacterium glutamicum* DSM 5715. Strain DSM 5715 is an
10 AEC-resistant lysine producer. The vector pMC1 cannot replicate independently in DSM5715 and is retained in the cell only if it has integrated into the chromosome of DSM 5715. Selection of clones with pMC1 integrated into the chromosome was carried out by plating out the
15 electroporation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which had been supplemented with 15 mg/l kanamycin. For detection of the integration, the internal *pgi* fragment (Example 5)
20 was labelled with the Dig hybridization kit from Boehringer Mannheim by the method of "The DIG System Users Guide for Filter Hybridization" of Boehringer Mannheim GmbH (Mannheim, Germany, 1993). Chromosomal DNA of a transformant was isolated by the method of Eikmanns et al.
25 (Microbiology 140: 1817 - 1828 (1994)) and in each case cleaved with the restriction enzymes SalI, SacI and HindIII. The fragments formed were separated by agarose gel electrophoresis and hybridized at 68°C with the Dig
hybridization kit from Boehringer. It was found in this way
30 that the plasmid pMC1 was inserted within the chromosomal *pgi* gene of strain DSM5715. The strain was called DSM5715::pMC1.

Example 7

Effect of over-expression of the zwf gene with simultaneous elimination of the pgi gene on the preparation of lysine

7.1 Preparation of the strain DSM5715::pMC1/pEC-T18mob2zwf

- 5 The vector pEC-T18mob2zwf mentioned in Example 1.2 was electroporated by the electroporation method of Tauch et al. (1994, FEMS Microbiological Letters, 123:343-347) in *Corynebacterium glutamicum* DSM 5715::pMC1. Selection for plasmid-carrying cells was made by plating out the
- 10 electroporation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), which had been supplemented with 15 mg/l kanamycin and with
- 15 5 mg/l tetracycline. Plasmid DNA was isolated from a transformant by conventional methods (Peters-Wendisch et al., 1998, Microbiology 144, 915-927) and checked by treatment with the restriction enzymes KpnI and SalI with subsequent agarose gel electrophoresis. The strain was called DSM5715::pMC1/pEC-T18mob2zwf.

20 7.2 Preparation of lysine

The *C. glutamicum* strain DSM5715::pMC1/pEC-T18mob2zwf obtained in Example 7.1 was cultured in a nutrient medium suitable for the production of lysine and the lysine content in the culture supernatant was determined.

- 25 For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with tetracycline (5 mg/l) and kanamycin (25 mg/l)) for 24 hours at 33°C. The cultures of the comparison strains were supplemented according to their resistance to antibiotics.
- 30 Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture.

Medium Cg III

NaCl 2.5 g/l

Bacto-Peptone 10 g/l

Bacto-Yeast extract 10 g/l

Glucose (autoclaved separately) 2% (w/v)

The pH was brought to pH 7.4

Tetracycline (5 mg/l) and kanamycin (5 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660 nm) of
5 the main culture was 0.1. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50 g/l
$(\text{NH}_4)_2\text{SO}_4$	25 g/l
KH_2PO_4	0.1 g/l
$\text{MgSO}_4 \cdot 7 \text{ H}_2\text{O}$	1.0 g/l
$\text{CaCl}_2 \cdot 2 \text{ H}_2\text{O}$	10 mg/l
$\text{FeSO}_4 \cdot 7 \text{ H}_2\text{O}$	10 mg/l
$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
L-Leucine (sterile-filtered)	0.1 g/l
CaCO_3	25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO_3 autoclaved in the dry state.

Culturing is carried out in a 10 ml volume in a 100 ml conical flask with baffles. Tetracycline (5 mg/l) and kanamycin (25 mg/l) were added. Culturing was carried out at 33°C and 80% atmospheric humidity.

After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-
5 BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in Table 3.

Table 3

Strain	OD	L-Lysine HCl g/l
DSM5715	7.3	14.3
DSM5715/pEC-T18mob2zwf	7.1	14.6
DSM5715::pMC1/ pECTmob2zwf	10.4	15.2

10 Example 8

Preparation of a genomic cosmid gene library from
Corynebacterium glutamicum ATCC 13032

Chromosomal DNA from Corynebacterium glutamicum ATCC 13032 was isolated as described by Tauch et al., (1995, Plasmid
15 33:168-179), and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product
20 Description SAP, Code no. 1758250). The DNA of the cosmid vector SuperCos1 (Wahl et al. (1987) Proceedings of the National Academy of Sciences USA 84:2160-2164), obtained from Stratagene (La Jolla, USA, Product Description SuperCos1 Cosmid Vektor Kit, Code no. 251301) was cleaved

with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline phosphatase. The cosmid DNA was then cleaved with the

5 restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032 DNA and the batch was treated with T4 DNA

10 ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04). The ligation mixture was then packed in phages with the aid of Gigapack II XL Packing Extracts (Stratagene, La Jolla, USA, Product Description Gigapack II XL Packing Extract, Code

15 no. 200217). For infection of the E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acid Res. 16:1563-1575) the cells were taken up in 10 mM MgSO₄ and mixed with an aliquot of the phage suspension. The infection and titering of the cosmid library were carried out as described by

20 Sambrook et al. (1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190) + 100 µg/ml ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

Example 9

25 Isolation and sequencing of the poxB gene

The cosmid DNA of an individual colony (Example 7) was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the

30 restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Product No.

35 1758250). After separation by gel electrophoresis, the

cosmid fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany). The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, Holland, Product Description Zero Background Cloning Kit, Product No. K2500-01), was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol Letters, 123:343-7) into the E. coli strain DH5 α MCR (Grant, 1990, Proceedings of the National Academy of Sciences U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 μ g/ml zeocin. The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain-stopping method of Sanger et al. (1977, Proceedings of the National Academies of Sciences U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No. Al24.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the

pZerol derivatives were assembled to a continuous contig. The computer-assisted coding region analysis were prepared with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231). Further analyses were carried out
5 with the "BLAST search program" (Altschul et al., 1997, Nucleic Acids Research, 25:3389-3402), against the non-redundant databank of the "National Center for Biotechnology Information" (NCBI, Bethesda, MD, USA).

The resulting nucleotide sequence is shown in SEQ ID No. 4.
10 Analysis of the nucleotide sequence showed an open reading frame of 1737 base pairs, which was called the poxB gene. The poxB gene codes for a polypeptide of 579 amino acids (SEQ ID NO. 5).

Example 10

15 Preparation of an integration vector for integration mutagenesis of the poxB gene

From the strain ATCC 13032, chromosomal DNA was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)). On the basis of the sequence of the poxB gene
20 known for *C. glutamicum* from Example 8, the following oligonucleotides were chosen for the polymerase chain reaction:

poxBint1:

5' TGC GAG ATG GTG AAT GGT GG 3'

25 poxBint2:

5' GCA TGA GGC AAC GCA TTA GC 3'

The primers shown were synthesized by MWG Biotech (Ebersberg, Germany) and the PCR reaction was carried out by the standard PCR method of Innis et al. (PCR protocols.
30 A guide to methods and applications, 1990, Academic Press) with Pwo-Polymerase from Boehringer. With the aid of the polymerase chain reaction, a DNA fragment approx. 0.9 kb in

size was isolated, this carrying an internal fragment of the poxB gene and being shown in SEQ ID No. 6.

The amplified DNA fragment was ligated with the TOPO TA Cloning Kit from Invitrogen Corporation (Carlsbad, CA, USA; Catalogue Number K4500-01) in the vector pCR2.1-TOPO (Mead at al. (1991) Bio/Technology 9:657-663). The E. coli strain DH5 α was then electroporated with the ligation batch (Hanahan, In: DNA cloning. A practical approach. Vol.I. IRL-Press, Oxford, Washington DC, USA, 1985). Selection for plasmid-carrying cells was made by plating out the transformation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), which had been supplemented with 25 mg/l kanamycin. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by restriction with the restriction enzyme EcoRI and subsequent agarose gel electrophoresis (0.8%). The plasmid was called pCR2.1poxBint (Figure 5).

Plasmid pCR2.1poxBint has been deposited in the form of the strain Escherichia coli DH5 α /pCR2.1poxBint as DSM 13114 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty.

Example 11

Integration mutagenesis of the poxB gene in the lysine producer DSM 5715

The vector pCR2.1poxBint mentioned in Example 10 was electroporated by the electroporation method of Tauch et al. (FEMS Microbiological Letters, 123:343-347 (1994)) in Corynebacterium glutamicum DSM 5715. Strain DSM 5715 is an AEC-resistant lysine producer. The vector pCR2.1poxBint

cannot replicate independently in DSM5715 and is retained in the cell only if it has integrated into the chromosome of DSM 5715. Selection of clones with pCR2.1poxBint integrated into the chromosome was carried out by plating out the electroporation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which had been supplemented with 15 mg/l kanamycin. For detection of the integration, the poxBint fragment was labelled with the Dig hybridization kit from Boehringer by the method of "The DIG System Users Guide for Filter Hybridization" of Boehringer Mannheim GmbH (Mannheim, Germany, 1993). Chromosomal DNA of a potential integrant was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)) and in each case cleaved with the restriction enzymes SalI, SacI and HindIII. The fragments formed were separated by agarose gel electrophoresis and hybridized at 68°C with the Dig hybridization kit from Boehringer. The plasmid pCR2.1poxBint mentioned in Example 9 had been inserted into the chromosome of DSM5715 within the chromosomal poxB gene. The strain was called DSM5715::pCR2.1poxBint.

Example 12

Effect of over-expression of the zwf gene with simultaneous elimination of the poxB gene on the preparation of lysine

12.1 Preparation of the strain DSM5715::pCR2.1poxBint/pEC-T18mob2zwf

The strain DSM5715::pCR2.1poxBint was transformed with the plasmid pEC-T18mob2zwf using the electroporation method described by Liebl et al., (FEMS Microbiology Letters, 53:299-303 (1989)). Selection of the transformants took place on LBHIS agar comprising 18.5 g/l brain-heart infusion broth, 0.5 M sorbitol, 5 g/l Bacto-tryptone, 2.5 g/l Bacto-yeast extract, 5 g/l NaCl and 18 g/l Bacto-

agar, which had been supplemented with 5 mg/l tetracycline and 25 mg/l kanamycin. Incubation was carried out for 2 days at 33°C.

Plasmid DNA was isolated in each case from a transformant
5 by conventional methods (Peters-Wendisch et al., 1998, Microbiology 144, 915-927), cleaved with the restriction endonucleases XbaI and KpnI, and the plasmid was checked by subsequent agarose gel electrophoresis. The strain obtained in this way was called DSM5715:pCR2.1poxBint/pEC-
10 T18mob2zwf.

12.2 Preparation of L-lysine

The *C. glutamicum* strain DSM5715::pCR2.1poxBint/pEC-T18mob2zwf obtained in Example 12.1 was cultured in a nutrient medium suitable for the production of lysine and
15 the lysine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with tetracycline (5 mg/l) and kanamycin (25 mg/l)) for 24 hours
20 at 33°C. The comparison strains were supplemented according to their resistance to antibiotics. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture.

Medium Cg III

NaCl	2.5 g/l
Bacto-Peptide	10 g/l
Bacto-Yeast extract	10 g/l
Glucose (autoclaved separately)	2% (w/v)

The pH was brought to pH 7.4

Tetracycline (5 mg/l) and kanamycin (25 mg/l) were added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660nm) of the main culture was 0.1. Medium MM was used for the main culture.

Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	58 g/l
$(\text{NH}_4)_2\text{SO}_4$	25 g/l
KH_2PO_4	0.1 g/l
$\text{MgSO}_4 \cdot 7 \text{ H}_2\text{O}$	1.0 g/l
$\text{CaCl}_2 \cdot 2 \text{ H}_2\text{O}$	10 mg/l
$\text{FeSO}_4 \cdot 7 \text{ H}_2\text{O}$	10 mg/l
$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
L-Leucine (sterile-filtered)	0.1 g/l
CaCO_3	25 g/l

The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO_3 autoclaved in the dry state.

- 5 Culturing is carried out in a 10 ml volume in a 100 ml conical flask with baffles. Tetracycline (5 mg/l) and kanamycin (25 mg/l) were added. Culturing was carried out at 33°C and 80% atmospheric humidity.

- 10 After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

- 15 The result of the experiment is shown in Table 4.

Table 4

Strain	OD	L-Lysine HCl g/l
DSM5715	10.8	16.0
DSM5715/pEC-T18mob2zwf	8.3	17.1
DSM5715::pCR2.1poxBint	7.1	16.7
DSM5715::pCR2.1poxBint/ pEC-Tmob2zwf	7.8	17.7

PCT


990239 BT

Original (for SUBMISSION) - printed on 03.07.2000 02:54:24 PM

0-1	Form - PCT/RO/134 (EASY) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared using	PCT-EASY Version 2.90 (updated 08.03.2000)
0-2	International Application No.	PCT/EP 00 / 0 6 3 0 3
0-3	Applicant's or agent's file reference	990239 BT

1	The Indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	page	10
1-2	line	24-29
1-3	Identification of Deposit	
1-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
1-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
1-3-3	Date of deposit	20 January 2000 (20.01.2000)
1-3-4	Accession Number	DSMZ 13244
1-4	Additional Indications	NONE
1-5	Designated States for Which Indications are Made	all designated States
1-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE

FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the international application: (yes or no)	yes
0-4-1	Authorized officer	 NATHALIE KUIPER

FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the International Bureau on:	
0-5-1	Authorized officer	

Patent claims

1. A process for the preparation of L-amino acids by
fermentation of coryneform bacteria
5 w h i c h c o m p r i s e s
 carrying out the following steps:
 - a) fermentation of the desired L-amino acid-
 producing bacteria in which at least the zwf gene
 is amplified,
 - 10 b) concentration of the L-amino acid in the medium
 or in the cells of the bacteria and
 - c) isolation of the L-amino acid produced.
2. The process as claimed in claim 1,
 w h e r e i n
15 bacteria in which further genes of the biosynthesis
 pathway of the desired L-amino acid are additionally
 amplified, in particular over-expressed, are employed.
3. The process as claimed in claim 1,
 w h e r e i n
20 coryneform bacteria which prepare L-threonine, L-lysine
 or L-tryptophan are used.
4. The process as claimed in claim 3,
 w h e r e i n
 coryneform bacteria which prepare L-lysine are used.
- 25 5. A process for the fermentative preparation of L-lysine
 as claimed in claim 2,
 w h e r e i n
 in the coryneform microorganisms which in particular
 already produce L-lysine, one or more genes chosen
30 from the group consisting of

- 5.1 the dapA gene which codes for dihydrodipicolinate synthase,
- 5.2 the lysC gene which codes for a feed back resistant aspartate kinase,
- 5 5.3 the gap gene which codes for glycerolaldehyde 3-phosphate dehydrogenase,
- 5.4 the pyc gene which codes for pyruvate carboxylase,
- 5.5 the tkt gene which codes for transketolase,
- 10 5.6 the gnd gene which codes for glucose 6-phosphate dehydrogenase,
- 5.7 the lysE gene which codes for lysine export,
- 5.8 the zwf gene,
- 5.9 the eno gene which codes for enolase
- 15 is or are amplified or over-expressed at the same time.
- 6. A process for the fermentative preparation of L-threonine as claimed in claim 2,
w h e r e i n
20 in the coryneform microorganisms which in particular already produce L-threonine, one or more genes chosen from the group consisting of
 - 6.1 the hom gene which codes for homoserine dehydrogenase or the hom^{dr} allele which codes for
25 a "feed back resistant" homoserine dehydrogenase,
 - 6.2 the gap gene which codes for glycerolaldehyde 3-phosphate dehydrogenase,


- 6.3 the *pyc* gene which codes for pyruvate carboxylase,
- 6.4 the *mgo* gene which codes for malate:quinone oxidoreductase,
- 5 6.5 the *tkt* gene which codes for transketolase,
- 6.6 the *zwf* gene which codes for glucose 6-phosphate dehydrogenase,
- 6.7 the *thrE* gene which codes for threonine export,
- 6.8 the *zwa1* gene,
- 10 6.9 the *eno* gene which codes for enolase
- is or are amplified, in particular over-expressed, at the same time.
7. The process as claimed in claim 2, where in
- 15 for the preparation of L-amino acids, in particular L-lysine or L-threonine, bacteria in which one or more genes chosen from the group consisting of,
- 7.1 the *pck* gene which codes for phosphoenol pyruvate carboxykinase,
- 20 7.2 the *pgi* gene which codes for glucose 6-phosphate isomerase
- 7.3 the *poxB* gene which codes for pyruvate oxidase or
- 7.4 the *zwa2* gene
- is or are attenuated at the same time, are fermented.
- 25 8. The process as claimed in claims 2 to 6, where in to achieve the amplification, the number of copies of

the genes or nucleotide sequences is increased by transformation of the microorganisms with plasmid vectors which carry these genes or nucleotide sequences.

- 5 9. The plasmid vector pEC-T18mob2 deposited under the designation DSM13244 in E.coli K-12 DH5 α , shown in Figure 2.
- 10 10. A coryneform microorganism, in particular of the genus *Corynebacterium*, transformed by the introduction of the plasmid vector as claimed in claim 9, which additionally contains the *zwf* gene.


1/5

Figure 1: Map of the plasmid pEC-T18mob2



2/5

Figure 2: Map of the plasmid pEC-T18mob2zwf



3/5

Figure 3






Figure 4:



5/5

Figure 5:



1

SEQUENCE PROTOCOL

<110> Degussa-Hüls AG
 National University of Ireland, Galway
 Forschungszentrum Jülich GmbH
 5 <120> Process for the fermentative preparation of
 L-amino acids using the zwf gene.
 <130> 990239BT
 10 <140>
 <141>
 <160> 6
 15 <170> PatentIn Ver. 2.1
 <210> 1
 <211> 2811
 20 <212> DNA
 <213> Corynebacterium glutamicum
 <220>
 <221> CDS
 25 <222> (373) .. (2022)
 <223> pgi
 <400> 1
 30 aaaacccgag gggcgaaaat tccaccctaa cttttttggg atcccccttt tccggggaat 60
 taattgggttt ggggtttcaat gggaaaaacgg gaaacaatgg gccaaagggt caaaaacccc 120
 aaaaggggggc cggtttcaaa ttcccaaaaa aaatggcaaa aaaggggggg ccaaaaccaa 180
 35 gttggccccc aaaccaccgg ggcaacggcc caccacaaaa ggggttgggt taaaggaagg 240
 acgcccacaaag taagcccgga atggcccacg ttcgaaaaag caggccccc aaacacgcac 300
 cttaaatttg tcgtgtttcc cactttgaac actcttcgat gcgcttgcc acaaaagcaa 360
 40 gctaacctga ag atg tta ttt aac gac aat aaa gga gtt ttc atg gcg gac 411
 Met Leu Phe Asn Asp Asn Lys Gly Val Phe Met Ala Asp
 1 5 10
 45 att tcg acc acc cag gtt tgg caa gac ctg acc gat cat tac tca aac 459
 Ile Ser Thr Thr Gln Val Trp Gln Asp Leu Thr Asp His Tyr Ser Asn
 15 20 25
 ttc cag gca acc act ctg cgt gaa ctt ttc aag gaa gaa aac cgc gcc 507
 50 Phe Gln Ala Thr Thr Leu Arg Glu Leu Phe Lys Glu Glu Asn Arg Ala
 30 35 40 45
 gag aag tac acc ttc tcc gcg gct ggc ctc cac gtc gac ctg tcg aag 555
 55 Glu Lys Tyr Thr Phe Ser Ala Ala Gly Leu His Val Asp Leu Ser Lys
 50 55 60
 aat ctg ctt gac gac gcc acc ctc acc aag ctc ctt gca ctg acc gaa 603

2

	Asn	Leu	Leu	Asp	Asp	Ala	Thr	Leu	Thr	Lys	Leu	Leu	Ala	Leu	Thr	Glu	
				65					70					75			
5	gaa	tct	ggc	ctt	cgc	gaa	cgc	att	gac	gcg	atg	ttt	gcc	ggt	gaa	cac	651
	Glu	Ser	Gly	Leu	Arg	Glu	Arg	Ile	Asp	Ala	Met	Phe	Ala	Gly	Glu	His	
			80					85					90				
10	ctc	aac	aac	acc	gaa	gac	cgc	gct	gtc	ctc	cac	acc	gcg	ctg	cgc	ctt	699
	Leu	Asn	Asn	Thr	Glu	Asp	Arg	Ala	Val	Leu	His	Thr	Ala	Leu	Arg	Leu	
		95					100					105					
15	cct	gcc	gaa	gct	gat	ctg	tca	gta	gat	ggc	caa	gat	gtt	gct	gct	gat	747
	Pro	Ala	Glu	Ala	Asp	Leu	Ser	Val	Asp	Gly	Gln	Asp	Val	Ala	Ala	Asp	
	110					115				120						125	
	gtc	cac	gaa	gtt	ttg	gga	cgc	atg	cgt	gac	ttc	gct	act	gcg	ctg	cgc	795
	Val	His	Glu	Val	Leu	Gly	Arg	Met	Arg	Asp	Phe	Ala	Thr	Ala	Leu	Arg	
					130					135					140		
20	tca	ggc	aac	tgg	ttg	gga	cac	acc	ggc	cac	acg	atc	aag	aag	atc	gtc	843
	Ser	Gly	Asn	Trp	Leu	Gly	His	Thr	Gly	His	Thr	Ile	Lys	Lys	Ile	Val	
				145					150						155		
25	aac	att	ggt	atc	ggt	ggc	tct	gac	ctc	gga	cca	gcc	atg	gct	acg	aag	891
	Asn	Ile	Gly	Ile	Gly	Gly	Ser	Asp	Leu	Gly	Pro	Ala	Met	Ala	Thr	Lys	
			160					165					170				
30	gct	ctg	cgt	gca	tac	gcg	acc	gct	ggt	atc	tca	gca	gaa	ttc	gtc	tcc	939
	Ala	Leu	Arg	Ala	Tyr	Ala	Thr	Ala	Gly	Ile	Ser	Ala	Glu	Phe	Val	Ser	
		175					180					185					
35	aac	gtc	gac	cca	gca	gac	ctc	gtt	tct	gtg	ttg	gaa	gac	ctc	gat	gca	987
	Asn	Val	Asp	Pro	Ala	Asp	Leu	Val	Ser	Val	Leu	Glu	Asp	Leu	Asp	Ala	
	190					195				200						205	
	gaa	tcc	aca	ttg	ttc	gtg	atc	gct	tcg	aaa	act	ttc	acc	acc	cag	gag	1035
	Glu	Ser	Thr	Leu	Phe	Val	Ile	Ala	Ser	Lys	Thr	Phe	Thr	Thr	Gln	Glu	
					210					215					220		
40	acg	ctg	tcc	aac	gct	cgt	gca	gct	cgt	gct	tgg	ctg	gta	gag	aag	ctc	1083
	Thr	Leu	Ser	Asn	Ala	Arg	Ala	Ala	Arg	Ala	Trp	Leu	Val	Glu	Lys	Leu	
				225					230					235			
45	ggt	gaa	gag	gct	gtc	gcg	aag	cac	ttc	gtc	gca	gtg	tcc	acc	aat	gct	1131
	Gly	Glu	Glu	Ala	Val	Ala	Lys	His	Phe	Val	Ala	Val	Ser	Thr	Asn	Ala	
			240					245					250				
50	gaa	aag	gtc	gca	gag	ttc	ggt	atc	gac	acg	gac	aac	atg	ttc	ggc	ttc	1179
	Glu	Lys	Val	Ala	Glu	Phe	Gly	Ile	Asp	Thr	Asp	Asn	Met	Phe	Gly	Phe	
		255					260					265					
55	tgg	gac	tgg	gtc	gga	ggt	cgt	tac	tcc	gtg	gac	tcc	gca	gtt	ggt	ctt	1227
	Trp	Asp	Trp	Val	Gly	Gly	Arg	Tyr	Ser	Val	Asp	Ser	Ala	Val	Gly	Leu	
	270					275					280					285	
	tcc	ctc	atg	gca	gtg	atc	ggc	cct	cgc	gac	ttc	atg	cgt	ttc	ctc	ggt	1275
	Ser	Leu	Met	Ala	Val	Ile	Gly	Pro	Arg	Asp	Phe	Met	Arg	Phe	Leu	Gly	
				290						295					300		

3

	gga ttc cac gcg atg gat gaa cac ttc cgc acc acc aag ttc gaa gag	1323
	Gly Phe His Ala Met Asp Glu His Phe Arg Thr Thr Lys Phe Glu Glu	
	305 310 315	
5	aac gtt cca atc ttg atg gct ctg ctc ggt gtc tgg tac tcc gat ttc	1371
	Asn Val Pro Ile Leu Met Ala Leu Leu Gly Val Trp Tyr Ser Asp Phe	
	320 325 330	
10	tat ggt gca gaa acc cac gct gtc cta cct tat tcc gag gat ctc agc	1419
	Tyr Gly Ala Glu Thr His Ala Val Leu Pro Tyr Ser Glu Asp Leu Ser	
	335 340 345	
15	cgt ttt gct gct tac ctc cag cag ctg acc atg gag acc aat ggc aag	1467
	Arg Phe Ala Ala Tyr Leu Gln Gln Leu Thr Met Glu Thr Asn Gly Lys	
	350 355 360 365	
20	tca gtc cac cgc gac ggc tcc cct gtt tcc act ggc act ggc gaa att	1515
	Ser Val His Arg Asp Gly Ser Pro Val Ser Thr Gly Thr Gly Glu Ile	
	370 375 380	
25	tac tgg ggt gag cct ggc aca aat ggc cag cac gct ttc ttc cag ctg	1563
	Tyr Trp Gly Glu Pro Gly Thr Asn Gly Gln His Ala Phe Phe Gln Leu	
	385 390 395	
30	atc cac cag ggc act cgc ctt gtt cca gct gat ttc att ggt ttc gct	1611
	Ile His Gln Gly Thr Arg Leu Val Pro Ala Asp Phe Ile Gly Phe Ala	
	400 405 410	
35	cgt cca aag cag gat ctt cct gcc ggt gag cgc acc atg cat gac ctt	1659
	Arg Pro Lys Gln Asp Leu Pro Ala Gly Glu Arg Thr Met His Asp Leu	
	415 420 425	
40	ttg atg agc aac ttc ttc gca cag acc aag gtt ttg gct ttc ggt aag	1707
	Leu Met Ser Asn Phe Phe Ala Gln Thr Lys Val Leu Ala Phe Gly Lys	
	430 435 440 445	
45	aac gct gaa gag atc gct gcg gaa ggt gtc gca cct gag ctg gtc aac	1755
	Asn Ala Glu Glu Ile Ala Ala Glu Gly Val Ala Pro Glu Leu Val Asn	
	450 455 460	
50	cac aag gtc gtg cca ggt aat cgc cca acc acc acc att ttg gcg gag	1803
	His Lys Val Val Pro Gly Asn Arg Pro Thr Thr Thr Ile Leu Ala Glu	
	465 470 475	
55	gaa ctt acc cct tct att ctc ggt gcg ttg atc gct ttg tac gaa cac	1851
	Glu Leu Thr Pro Ser Ile Leu Gly Ala Leu Ile Ala Leu Tyr Glu His	
	480 485 490	
60	acc gtg atg gtt cag ggc gtg att tgg gac atc aac tcc ttc gac caa	1899
	Thr Val Met Val Gln Gly Val Ile Trp Asp Ile Asn Ser Phe Asp Gln	
	495 500 505	
65	tgg ggt gtt gaa ctg ggc aaa cag cag gca aat gac ctc gct ccg gct	1947
	Trp Gly Val Glu Leu Gly Lys Gln Gln Ala Asn Asp Leu Ala Pro Ala	
	510 515 520 525	
70	gtc tct ggt gaa gag gat gtt gac tcg gga gat tct tcc act gat tca	1995
	Val Ser Gly Glu Asp Val Asp Ser Gly Asp Ser Ser Thr Asp Ser	
	530 535 540	

ctg att aag tgg tac cgc gca aat agg tagtcgcttg cttatagggt 2042
 Leu Ile Lys Trp Tyr Arg Ala Asn Arg
 545 550

5

caggggacgtg aagaatcctc gcctcatagc actggccgct atcatcctga cctcgttcaa 2102
 tctgcgaaca gctattactg ctttagctcc gctggtttct gagattcggg atgatttagg 2162
 10 ggttagtgct tctcttattg gtgtgttggt catgatcccg actgctatgt tcgcggttgc 2222
 tgcgtttgcg cttccgctgt tgaagaggaa gtccactact tcccaactgt tgatgtttgc 2282
 catgctgttg actgctgccg gtcagattat tcgtgtcgtt ggacctgctt cgctgttgat 2342
 15 ggtcgggtact gtgttcgcga tgtttgcgat cggagttacc aatgtgttgc ttccgattgc 2402
 tgttagggag tattttccgc gtcacgtcgg tggaatgtcg acaacttatc tgggtgcgtt 2462
 20 ccagattgtt caggcacttg ctccgacgct tgccgtgccg atttctcagt gggctacaca 2522
 tgtgggggttg accggttgga ggggtgcgct cggttcgtgg gcgctgctgg ggttggttgc 2582
 ggcgatttcg tggattccgc tgttgagttt gcagggtgcc aggggtgttg cggcgccgctc 2642
 25 gaaggtttct cttcctgtgt ggaagtcttc ggttggtgtg gggctcgggt tgatgtttgg 2702
 gtttacttcg tttgcgacgt atatcctcat gggttttatg ccgcagatgg taggtgatcc 2762
 30 aaagaattca aaaagcttct cgagagtact tctagagcgg ccgcgggcc 2811

<210> 2
 <211> 550
 35 <212> PRT
 <213> Corynebacterium glutamicum

<400> 2
 Met Leu Phe Asn Asp Asn Lys Gly Val Phe Met Ala Asp Ile Ser Thr
 40 1 5 10 15
 Thr Gln Val Trp Gln Asp Leu Thr Asp His Tyr Ser Asn Phe Gln Ala
 20 25 30
 45 Thr Thr Leu Arg Glu Leu Phe Lys Glu Glu Asn Arg Ala Glu Lys Tyr
 35 40 45
 Thr Phe Ser Ala Ala Gly Leu His Val Asp Leu Ser Lys Asn Leu Leu
 50 55 60
 50 Asp Asp Ala Thr Leu Thr Lys Leu Leu Ala Leu Thr Glu Glu Ser Gly
 65 70 75 80
 Leu Arg Glu Arg Ile Asp Ala Met Phe Ala Gly Glu His Leu Asn Asn
 55 85 90 95
 Thr Glu Asp Arg Ala Val Leu His Thr Ala Leu Arg Leu Pro Ala Glu
 100 105 110

5

Ala Asp Leu Ser Val Asp Gly Gln Asp Val Ala Ala Asp Val His Glu
 115 120 125
 Val Leu Gly Arg Met Arg Asp Phe Ala Thr Ala Leu Arg Ser Gly Asn
 5 130 135 140
 Trp Leu Gly His Thr Gly His Thr Ile Lys Lys Ile Val Asn Ile Gly
 145 150 155 160
 10 Ile Gly Gly Ser Asp Leu Gly Pro Ala Met Ala Thr Lys Ala Leu Arg
 165 170 175
 Ala Tyr Ala Thr Ala Gly Ile Ser Ala Glu Phe Val Ser Asn Val Asp
 180 185 190
 15 Pro Ala Asp Leu Val Ser Val Leu Glu Asp Leu Asp Ala Glu Ser Thr
 195 200 205
 20 Leu Phe Val Ile Ala Ser Lys Thr Phe Thr Thr Gln Glu Thr Leu Ser
 210 215 220
 Asn Ala Arg Ala Ala Arg Ala Trp Leu Val Glu Lys Leu Gly Glu Glu
 225 230 235 240
 25 Ala Val Ala Lys His Phe Val Ala Val Ser Thr Asn Ala Glu Lys Val
 245 250 255
 Ala Glu Phe Gly Ile Asp Thr Asp Asn Met Phe Gly Phe Trp Asp Trp
 260 265 270
 30 Val Gly Gly Arg Tyr Ser Val Asp Ser Ala Val Gly Leu Ser Leu Met
 275 280 285
 35 Ala Val Ile Gly Pro Arg Asp Phe Met Arg Phe Leu Gly Gly Phe His
 290 295 300
 Ala Met Asp Glu His Phe Arg Thr Thr Lys Phe Glu Glu Asn Val Pro
 305 310 315 320
 40 Ile Leu Met Ala Leu Leu Gly Val Trp Tyr Ser Asp Phe Tyr Gly Ala
 325 330 335
 Glu Thr His Ala Val Leu Pro Tyr Ser Glu Asp Leu Ser Arg Phe Ala
 340 345 350
 45 Ala Tyr Leu Gln Gln Leu Thr Met Glu Thr Asn Gly Lys Ser Val His
 355 360 365
 50 Arg Asp Gly Ser Pro Val Ser Thr Gly Thr Gly Glu Ile Tyr Trp Gly
 370 375 380
 Glu Pro Gly Thr Asn Gly Gln His Ala Phe Phe Gln Leu Ile His Gln
 385 390 395 400
 55 Gly Thr Arg Leu Val Pro Ala Asp Phe Ile Gly Phe Ala Arg Pro Lys
 405 410 415
 Gln Asp Leu Pro Ala Gly Glu Arg Thr Met His Asp Leu Leu Met Ser
 420 425 430

Asn Phe Phe Ala Gln Thr Lys Val Leu Ala Phe Gly Lys Asn Ala Glu
 435 440 445

5 Glu Ile Ala Ala Glu Gly Val Ala Pro Glu Leu Val Asn His Lys Val
 450 455 460

Val Pro Gly Asn Arg Pro Thr Thr Thr Ile Leu Ala Glu Glu Leu Thr
 10 465 470 475 480

Pro Ser Ile Leu Gly Ala Leu Ile Ala Leu Tyr Glu His Thr Val Met
 485 490 495

Val Gln Gly Val Ile Trp Asp Ile Asn Ser Phe Asp Gln Trp Gly Val
 15 500 505 510

Glu Leu Gly Lys Gln Gln Ala Asn Asp Leu Ala Pro Ala Val Ser Gly
 515 520 525

20 Glu Glu Asp Val Asp Ser Gly Asp Ser Ser Thr Asp Ser Leu Ile Lys
 530 535 540

Trp Tyr Arg Ala Asn Arg
 25 545 550

<210> 3
 <211> 462
 30 <212> DNA
 <213> Corynebacterium glutamicum

<400> 3
 35 atggagacca atggcaagtc agtccaccgc gacggctccc ctgtttccac tggcactggc 60
 gaaatttact ggggtgagcc tggcacaaat ggccagcacg ctttcttcca gctgatccac 120
 cagggcactc gccttgttcc agctgatttc attggtttcg ctctccaaa gcaggatctt 180
 cctgccggtg agcgacccat gcatgacctt ttgatgagca acttcttcgc acagaccaag 240
 gttttggtt tcggttaagaa cgctgaagag atcgtgctcg aaggtgtcgc acctgagctg 300
 gtcaaccaca aggtcgtgcc aggtaatcgc ccaaccacca ccattttggc ggaggaactt 360
 40 accccttcta ttctcgtgac gttgatcgct ttgtacgaac acaccgtgat ggttcagggc 420
 gtgatttggg acatcaactc cttcgaccaaa tggggcgtgg aa 462

<210> 4
 45 <211> 2160
 <212> DNA
 <213> Corynebacterium glutamicum

<220>
 50 <221> CDS
 <222> (327)..(2063)
 <223> poxB

<400> 4
 55 ttagaggcga ttctgtgagg tcactttttg tggggctcgg gtctaaattt ggccagtttt 60
 cgaggcgacc agacaggcgt gccacgatg tttaaatagg cgatcgggtg gcatctgtgt 120
 ttggtttcga cgggctgaaa ccaaaccaga ctgcccagca acgacggaaa tcccaaaagt 180

	205	210	215	
5	cag gtg ttg gag ttg gcg gag aag att aaa tca ccg atc ggg cat gcg Gln Val Leu Glu Leu Ala Glu Lys Ile Lys Ser Pro Ile Gly His Ala 220 225 230	1025		
10	ctg ggt ggt aag cag tac atc cag cat gag aat ccg ttt gag gtc ggc Leu Gly Gly Lys Gln Tyr Ile Gln His Glu Asn Pro Phe Glu Val Gly 235 240 245	1073		
15	atg tct ggc ctg ctt ggt tac ggc gcc tgc gtg gat gcg tcc aat gag Met Ser Gly Leu Leu Gly Tyr Gly Ala Cys Val Asp Ala Ser Asn Glu 250 255 260 265	1121		
20	gcg gat ctg ctg att cta ttg ggt acg gat ttc cct tat tct gat ttc Ala Asp Leu Leu Ile Leu Leu Gly Thr Asp Phe Pro Tyr Ser Asp Phe 270 275 280	1169		
25	ctt cct aaa gac aac gtt gcc cag gtg gat atc aac ggt gcg cac att Leu Pro Lys Asp Asn Val Ala Gln Val Asp Ile Asn Gly Ala His Ile 285 290 295	1217		
30	ggt cga cgt acc acg gtg aag tat ccg gtg acc ggt gat gtt gct gca Gly Arg Arg Thr Thr Val Lys Tyr Pro Val Thr Gly Asp Val Ala Ala 300 305 310	1265		
35	aca atc gaa aat att ttg cct cat gtg aag gaa aaa aca gat cgt tcc Thr Ile Glu Asn Ile Leu Pro His Val Lys Glu Lys Thr Asp Arg Ser 315 320 325	1313		
40	ttc ctt gat cgg atg ctc aag gca cac gag cgt aag ttg agc tcg gtg Phe Leu Asp Arg Met Leu Lys Ala His Glu Arg Lys Leu Ser Ser Val 330 335 340 345	1361		
45	gta gag acg tac aca cat aac gtc gag aag cat gtg cct att cac cct Val Glu Thr Tyr Thr His Asn Val Glu Lys His Val Pro Ile His Pro 350 355 360	1409		
50	gaa tac gtt gcc tct att ttg aac gag ctg gcg gat aag gat gcg gtg Glu Tyr Val Ala Ser Ile Leu Asn Glu Leu Ala Asp Lys Asp Ala Val 365 370 375	1457		
55	ttt act gtg gat acc ggc atg tgc aat gtg tgg cat gcg agg tac atc Phe Thr Val Asp Thr Gly Met Cys Asn Val Trp His Ala Arg Tyr Ile 380 385 390	1505		
60	gag aat ccg gag gga acg cgc gac ttt gtg ggt tca ttc cgc cac ggc Glu Asn Pro Glu Gly Thr Arg Asp Phe Val Gly Ser Phe Arg His Gly 395 400 405	1553		
65	acg atg gct aat gcg ttg cct cat gcg att ggt gcg caa agt gtt gat Thr Met Ala Asn Ala Leu Pro His Ala Ile Gly Ala Gln Ser Val Asp 410 415 420 425	1601		
70	cga aac cgc cag gtg atc gcg atg tgt ggc gat ggt ggt ttg ggc atg Arg Asn Arg Gln Val Ile Ala Met Cys Gly Asp Gly Gly Leu Gly Met 430 435 440	1649		
75	ctg ctg ggt gag ctt ctg acc gtt aag ctg cac caa ctt ccg ctg aag	1697		

9

Leu Leu Gly Glu Leu Leu Thr Val Lys Leu His Gln Leu Pro Leu Lys
 445 450 455
 5 gct gtg gtg ttt aac aac agt tct ttg ggc atg gtg aag ttg gag atg 1745
 Ala Val Val Phe Asn Asn Ser Ser Leu Gly Met Val Lys Leu Glu Met
 460 465 470
 10 ctc gtg gag gga cag cca gaa ttt ggt act gac cat gag gaa gtg aat 1793
 Leu Val Glu Gly Gln Pro Glu Phe Gly Thr Asp His Glu Glu Val Asn
 475 480 485
 15 ttc gca gag att gcg gcg gct gcg ggt atc aaa tcg gta cgc atc acc 1841
 Phe Ala Glu Ile Ala Ala Ala Gly Ile Lys Ser Val Arg Ile Thr
 490 495 500 505
 gat ccg aag aaa gtt cgc gag cag cta gct gag gca ttg gca tat cct 1889
 Asp Pro Lys Lys Val Arg Glu Gln Leu Ala Glu Ala Leu Ala Tyr Pro
 510 515 520
 20 gga cct gta ctg atc gat atc gtc acg gat cct aat gcg ctg tcg atc 1937
 Gly Pro Val Leu Ile Asp Ile Val Thr Asp Pro Asn Ala Leu Ser Ile
 525 530 535
 25 cca cca acc atc acg tgg gaa cag gtc atg gga ttc agc aag gcg gcc 1985
 Pro Pro Thr Ile Thr Trp Glu Gln Val Met Gly Phe Ser Lys Ala Ala
 540 545 550
 30 acc cga acc gtc ttt ggt gga gga gta gga gcg atg atc gat ctg gcc 2033
 Thr Arg Thr Val Phe Gly Gly Gly Val Gly Ala Met Ile Asp Leu Ala
 555 560 565
 35 cgt tcg aac ata agg aat att cct act cca tgatgattga tacacctgct 2083
 Arg Ser Asn Ile Arg Asn Ile Pro Thr Pro
 570 575
 gttctcattg accgcgagcg cttaactgcc aacatttcca ggatggcagc tcacgccggt 2143
 gcccatgaga ttgccct 2160
 40 <210> 5
 <211> 579
 <212> PRT
 <213> Corynebacterium glutamicum
 45 <400> 5
 Met Ala His Ser Tyr Ala Glu Gln Leu Ile Asp Thr Leu Glu Ala Gln
 1 5 10 15
 50 Gly Val Lys Arg Ile Tyr Gly Leu Val Gly Asp Ser Leu Asn Pro Ile
 20 25 30
 Val Asp Ala Val Arg Gln Ser Asp Ile Glu Trp Val His Val Arg Asn
 35 40 45
 55 Glu Glu Ala Ala Ala Phe Ala Ala Gly Ala Glu Ser Leu Ile Thr Gly
 50 55 60
 Glu Leu Ala Val Cys Ala Ala Ser Cys Gly Pro Gly Asn Thr His Leu

	65				70				75				80			
5	Ile	Gln	Gly	Leu	Tyr	Asp	Ser	His	Arg	Asn	Gly	Ala	Lys	Val	Leu	Ala
					85					90					95	
	Ile	Ala	Ser	His	Ile	Pro	Ser	Ala	Gln	Ile	Gly	Ser	Thr	Phe	Phe	Gln
				100					105					110		
10	Glu	Thr	His	Pro	Glu	Ile	Leu	Phe	Lys	Glu	Cys	Ser	Gly	Tyr	Cys	Glu
			115					120					125			
	Met	Val	Asn	Gly	Gly	Glu	Gln	Gly	Glu	Arg	Ile	Leu	His	His	Ala	Ile
		130					135					140				
15	Gln	Ser	Thr	Met	Ala	Gly	Lys	Gly	Val	Ser	Val	Val	Val	Ile	Pro	Gly
	145					150					155					160
	Asp	Ile	Ala	Lys	Glu	Asp	Ala	Gly	Asp	Gly	Thr	Tyr	Ser	Asn	Ser	Thr
					165					170					175	
20	Ile	Ser	Ser	Gly	Thr	Pro	Val	Val	Phe	Pro	Asp	Pro	Thr	Glu	Ala	Ala
				180					185					190		
	Ala	Leu	Val	Glu	Ala	Ile	Asn	Asn	Ala	Lys	Ser	Val	Thr	Leu	Phe	Cys
			195					200					205			
	Gly	Ala	Gly	Val	Lys	Asn	Ala	Arg	Ala	Gln	Val	Leu	Glu	Leu	Ala	Glu
		210					215					220				
30	Lys	Ile	Lys	Ser	Pro	Ile	Gly	His	Ala	Leu	Gly	Gly	Lys	Gln	Tyr	Ile
	225					230					235					240
	Gln	His	Glu	Asn	Pro	Phe	Glu	Val	Gly	Met	Ser	Gly	Leu	Leu	Gly	Tyr
				245						250					255	
35	Gly	Ala	Cys	Val	Asp	Ala	Ser	Asn	Glu	Ala	Asp	Leu	Leu	Ile	Leu	Leu
				260					265					270		
	Gly	Thr	Asp	Phe	Pro	Tyr	Ser	Asp	Phe	Leu	Pro	Lys	Asp	Asn	Val	Ala
			275					280					285			
	Gln	Val	Asp	Ile	Asn	Gly	Ala	His	Ile	Gly	Arg	Arg	Thr	Thr	Val	Lys
		290					295					300				
45	Tyr	Pro	Val	Thr	Gly	Asp	Val	Ala	Ala	Thr	Ile	Glu	Asn	Ile	Leu	Pro
	305					310					315					320
	His	Val	Lys	Glu	Lys	Thr	Asp	Arg	Ser	Phe	Leu	Asp	Arg	Met	Leu	Lys
					325					330					335	
50	Ala	His	Glu	Arg	Lys	Leu	Ser	Ser	Val	Val	Glu	Thr	Tyr	Thr	His	Asn
				340					345					350		
	Val	Glu	Lys	His	Val	Pro	Ile	His	Pro	Glu	Tyr	Val	Ala	Ser	Ile	Leu
			355					360					365			
55	Asn	Glu	Leu	Ala	Asp	Lys	Asp	Ala	Val	Phe	Thr	Val	Asp	Thr	Gly	Met
		370					375						380			

11

Cys Asn Val Trp His Ala Arg Tyr Ile Glu Asn Pro Glu Gly Thr Arg
 385 390 395 400
 5 Asp Phe Val Gly Ser Phe Arg His Gly Thr Met Ala Asn Ala Leu Pro
 405 410 415
 His Ala Ile Gly Ala Gln Ser Val Asp Arg Asn Arg Gln Val Ile Ala
 420 425 430
 10 Met Cys Gly Asp Gly Gly Leu Gly Met Leu Leu Gly Glu Leu Leu Thr
 435 440 445
 Val Lys Leu His Gln Leu Pro Leu Lys Ala Val Val Phe Asn Asn Ser
 450 455 460
 15 Ser Leu Gly Met Val Lys Leu Glu Met Leu Val Glu Gly Gln Pro Glu
 465 470 475 480
 Phe Gly Thr Asp His Glu Glu Val Asn Phe Ala Glu Ile Ala Ala Ala
 485 490 495
 Ala Gly Ile Lys Ser Val Arg Ile Thr Asp Pro Lys Lys Val Arg Glu
 500 505 510
 25 Gln Leu Ala Glu Ala Leu Ala Tyr Pro Gly Pro Val Leu Ile Asp Ile
 515 520 525
 Val Thr Asp Pro Asn Ala Leu Ser Ile Pro Pro Thr Ile Thr Trp Glu
 530 535 540
 30 Gln Val Met Gly Phe Ser Lys Ala Ala Thr Arg Thr Val Phe Gly Gly
 545 550 555 560
 Gly Val Gly Ala Met Ile Asp Leu Ala Arg Ser Asn Ile Arg Asn Ile
 565 570 575
 35 Pro Thr Pro
 40
 <210> 6
 <211> 875
 <212> DNA
 45 <213> *Corynebacterium glutamicum*
 <400> 6
 50 tgcgagatgg tgaatggtgg tgagcagggg gaacgcattt tgcatacagc gattcagtc 60
 accatggcgg gtaaagggtg gtcggtggta gtgattcctg gtgatatcgc taaggaagac 120
 gcaggtgacg gtacttattc caattocact atttcttctg gcactcctgt ggtgttccc 180
 gatcctactg aggctgcagc gctggtggag gcgattaaca acgctaagtc tgtcactttg 240
 ttctgcggtg cgggcgtgaa gaatgctcgc gcgcaggtgt tggagttggc ggagaagatt 300
 aaatcaccga tcgggcatgc gctgggtggg aagcagtaca tccagcatga gaatccgttt 360
 gaggtcggca tgtctggcct gcttggttac ggcgcctgcg tggatgcgtc caatgaggcg 420
 55 gatctgctga ttctattggg tacggatttc ccttattctg atttccttcc taaagacaac 480
 gttgcccgagg tggatatcaa cggtgccgac attggtcgac gtaccacggg gaagtatccg 540
 gtgaccgggtg atgttgctgc aacaatcgaa aatattttgc ctcatgtgaa ggaaaaaaca 600
 gatcggttct tccttgatcg gatgctcaag gcacacgagc gtaagttgag ctccggtggt 660
 gagacgtaca cacataacgt cgagaagcat gtgcctattc accctgaata cgttgccctc 720

12

```
at t t t g a a c g   a g c t g g c g g a   t a a g g a t g c g   g t g t t t a c t g   t g g a t a c c g g   c a t g t g c a a t   780
g t g t g g c a t g   c g a g g t a c a t   c g a g a a t c c g   g a g g g a a c g c   g c g a c t t t g t   g g g t t c a t t c   840
c g c c a c g g c a   c g a t g g c t a a   t g c g t t g c c t   c a t g c                                     875
```

5

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/06303

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/53 C12N15/77 C12P13/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12P C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, MEDLINE, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>JP 09 224661 A (MITSUBISHI CHEM CORP) 2 September 1997 (1997-09-02) cited in the application the whole document & DATABASE WPI Section Ch, Week 199745 Derwent Publications Ltd., London, GB; Class B04, AN 1997-484096 HATAKEYAMA KAZUHISA ET AL.: "Glucose-6-phosphate dehydrogenase " & JP 09 224661 A (MITSUBISHI CHEM CORP), 2 September 1997 (1997-09-02) abstract</p> <p style="text-align: center;">--- -/--</p>	9, 10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *A* document member of the same patent family

Date of the actual completion of the international search

13 February 2001

Date of mailing of the international search report

21/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Montero Lopez, B

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 00/06303

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>L. EGGELING ET AL.: "L-Glutamate and L-Lysine: traditional products with impetuous developments"</p> <p>APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, vol. 52, August 1999 (1999-08), pages 146-153, XP000979507</p> <p>the whole document</p> <p>-----</p>	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/06303

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 9224661 A	02-09-1997	NONE	